

## Corydoras size group 'B'

Over 57 mm SL. (females)

No	Name	C No	M	F	Comments
01	<i>C. acutus</i>		65	70	
02	<i>C. adolfoi</i>		55	60	
03	<i>C. aeneus</i> (albino)		65	70	
04	<i>C. aeneus</i>		65	75	
05	<i>C. agassizii</i>		65	70	
06	<i>C. amandajanea</i>		60	65	
07	<i>C. amapaensis</i>		60	70	
08	<i>C. ambiacus</i>		60	65	
09	<i>C. amphibelus</i>		55	60	No way to id, Type material lost.
10	<i>C. approuaguensis</i>		65	70	
11	<i>C. araguaiaensis</i>		55	60	
12	<i>C. areio</i>		60	65	
13	<i>C. armatus</i>		60	65	
14	<i>C. bifasciatus</i>		55	60	
15	<i>C. blochi</i>	C32	55	60	Formerly <i>C. blochi blochi</i>
16	<i>C. brevirostris</i>		55	60	
17	<i>C. cervinus</i>		60	70	
18	<i>C. concolor</i>		55	60	
19	<i>C. condiscipulus</i>		60	65	
20	<i>C. coriatae</i>		60	65	
21	<i>C. crimmeni</i>		60	65	
22	<i>C. crypticus</i>		60	65	
23	<i>C. davidsandsi</i>		55	60	
24	<i>C. delphax</i>		65	70	
25	<i>C. ellisae</i>		60	65	
26	<i>C. ephippifer</i>		60	65	
27	<i>C. evelynae</i>		55	60	
28	<i>C. fowleri</i>		65	70	
29	<i>C. geoffroy</i>		65	70	Formerly <i>C. octocirrus</i>
30	<i>C. geryi</i>		65	70	Formerly <i>C. bolivianus</i> (bars in caudal)
31	<i>C. gomezi</i>		55	60	
32	<i>C. gossei</i>		55	60	
33	<i>C. haraldschultzi</i>		80	85	
34	<i>C. heteromorphus</i>		60	65	
35	<i>C. imitator</i>		60	65	
36	<i>C. inolicana</i>	C1	70	80	
37	<i>C. isbueckeri</i>	C50	65	70	New species Knaack, 2004
38	<i>C. lamberti</i>		55	60	
39	<i>C. leopardus</i>		70	75	
40	<i>C. melanotaenia</i>		55	60	
41	<i>C. narcissus</i>		70	75	
42	<i>C. nattereri</i>		60	65	
43	<i>C. negro</i>		60	65	New species Knaack, 2004
44	<i>C. noelkempffi</i>	C58	60	65	New species Knaack, 2004
45	<i>C. orcesi</i>		65	70	
46	<i>C. ornatus</i>		65	70	
47	<i>C. orphnopterus</i>		65	70	
48	<i>C. ourastigma</i>		60	65	
49	<i>C. oxyrhynchus</i>		65	70	
50	<i>C. paleatus</i>		65	70	
51	<i>C. paleatus</i> (Albino)		60	65	
52	<i>C. pantanalensis</i>	C5	75	80	Sim. <i>C. geryi</i> but no bars in caudal
53	<i>C. parallelus</i>	C2	55	60	

54	<i>C. pastazensis</i>		65	70	
55	<i>C. pinheiroi</i>	C25	60	65	
56	<i>C. pulcher</i>		60	65	
57	<i>C. reticulatus</i>		60	65	
58	<i>C. robineae</i>		65	70	
59	<i>C. robustus</i>		85	90	
60	<i>C. saramaccensis</i>	C23	65	70	
61	<i>C. sararensis</i>		60	65	
62	<i>C. schultzi</i>		65	70	Not scientifically proved <i>C. cf aeneus</i> ***
63	<i>C. schwartzi</i>		55	60	
64	<i>C. semiaquilus</i>		75	80	
65	<i>C. septentrionalis</i>		50	60	
66	<i>C. serratus</i>		60	65	
67	<i>C. seussi</i>	C27	65	70	
68	<i>C. similis</i>		55	60	
69	<i>C. simulatus</i>		65	70	
70	<i>C. sodalis</i>		55	60	
71	<i>C. solox</i>		65	70	
72	<i>C. spectabilis</i>		65	70	
73	<i>C. stenocephalus</i>	C17	65	70	
74	<i>C. sterbai</i>		60	65	
75	<i>C. sychri</i>		60	65	
76	<i>C. treitlii</i>		65	70	
77	<i>C. virginiae</i>	C4	55	60	
78	<i>C. vittatus</i>		55	60	Formerly <i>C. blochi vittatus</i>
79	<i>C. zygatus</i>		65	70	

79 species.

Note: - There are no longer any sub-species within the family Corydoradinae (Isbrücker 2001).

There are three species scientifically regarded as synonyms of *C. aeneus*, but they are considerably different in looks or size.

<i>C. macrosteus</i>	70	75	A species from Argentina.
<i>C. schultzi</i>	65	70	has the gold shoulder flash which extends along the upper back.
<i>C. venezuelanus</i>	50	55	A small rounded species with dark ovate central body blotch.